



Crystal structure of *trans*-diaquabis(4-cyanobenzoato- κO)bis-(nicotinamide- κN^1)cobalt(II)

Gülçin Şefiye Aşkın, Hacali Necefoğlu, Gamze Yılmaz Nayir, Raziye Çatak
Çelik and Tuncer Hökelek

Acta Cryst. (2015). **E71**, 561–563



IUCr Journals

CRYSTALLOGRAPHY JOURNALS ONLINE

This open-access article is distributed under the terms of the Creative Commons Attribution Licence
<http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and
reproduction in any medium, provided the original authors and source are cited.



Received 22 April 2015
Accepted 27 April 2015

Edited by M. Weil, Vienna University of
Technology, Austria

Keywords: crystal structure; cobalt(II); transition metal complexes of benzoic acid and nicotinamide derivatives

CCDC reference: 1061935

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structure of *trans*-diaquabis(4-cyano-benzoato- κO)bis(nicotinamide- κN^1)cobalt(II)



Gülçin Şefiye Aşkın,^a Hacali Necefoğlu,^{b,c} Gamze Yılmaz Nayir,^b Raziye Çatak Çelik^d and Tuncer Hökelek^{a*}

^aDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey, ^bDepartment of Chemistry, Kafkas University, 36100 Kars, Turkey, ^cInternational Scientific Research Centre, Baku State University, 1148 Baku, Azerbaijan, and ^dScientific and Technological Application and Research Center, Aksaray University, 68100 Aksaray, Turkey.
*Correspondence e-mail: merzifon@hacettepe.edu.tr


In the title complex, $[Co(C_8H_4NO_2)_2(C_6H_6N_2O)_2(H_2O)_2]$, the Co^{II} atom is located on an inversion centre and is coordinated by two 4-cyanobenzoate (CNB) anions, two nicotinamide (NA) ligands and two water molecules. The four O atoms in the equatorial plane form a slightly distorted square-planar arrangement, while the slightly distorted octahedral coordination sphere is completed by the two N atoms of the NA ligands in the axial positions. The dihedral angle between the carboxylate group and the adjacent benzene ring is 22.11 (15) $^\circ$, while the pyridine and benzene rings are oriented at a dihedral angle of 89.98 (5) $^\circ$. In the crystal, intermolecular N—H···O and O—H···O hydrogen bonds link the molecules, enclosing $R_2^2(8)$ and $R_4^4(8)$ ring motifs, forming layers parallel to (100). The layers are linked via C—H···O and C—H···N hydrogen bonds, resulting in a three-dimensional network. A weak C—H··· π interaction is also observed.

1. Chemical context

Nicotinamide (NA) is one form of niacin. A deficiency of this vitamin leads to loss of copper from the body, known as pellagra disease. Victims of pellagra show unusually high serum and urinary copper levels (Krishnamachari, 1974). The nicotinic acid derivative *N,N*-diethylnicotinamide (DENA) is an important respiratory stimulant (Bigoli *et al.*, 1972). Transition metal complexes with biochemical-relevant molecules show interesting physical and/or chemical properties, through which they may find applications in biological systems (Antolini *et al.*, 1982). Some benzoic acid derivatives, such as 4-aminobenzoic acid, have been extensively reported in coordination chemistry, as bifunctional organic ligands, due to the varieties of their coordination modes (Chen & Chen, 2002; Amiraslanov *et al.*, 1979; Hauptmann *et al.*, 2000).



The structure–function–coordination relationships of the arylcarboxylate ion in Zn^{II} complexes of benzoic acid

**Figure 1**

The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Symmetry-related atoms are defined by symmetry code $-x + 1, -y + 1, -z + 1$.

derivatives change depending on the nature and position of the substituted groups on the benzene ring, the nature of the additional ligand molecule or solvent, and the pH and temperature of synthesis (Shnulin *et al.*, 1981; Nadzhafov *et al.*, 1981; Antsyshkina *et al.*, 1980; Adiwidjaja *et al.*, 1978). When pyridine and its derivatives are used instead of water molecules, the structure is completely different (Catterick *et al.*, 1974). In this context, we synthesized the Co^{II}-containing title compound, *trans*-diaqua^{bis}(4-cyanobenzoato- κO)bis(nicotinamide- κN^1)cobalt(II), [Co(C₈H₄O₂N)₂(C₆H₆N₂O)₂(H₂O)₂], and report herein its crystal structure.

2. Structural commentary

In the mononuclear title complex, the Co^{II} atom is located on an inversion centre and is coordinated by two 4-cyanobenzoate (CNB) anions, two nicotinamide (NA) ligands and two water molecules, with all ligands coordinating in a monodentate manner (Fig. 1).

The two symmetry-related carboxylate O atoms (O₂ and O₂ⁱ) and the two symmetry-related water O atoms (O₄ and O₄ⁱ) form a slightly distorted square-planar arrangement, while the slightly distorted octahedral coordination sphere is completed by the two symmetry-related N atoms (N₂ and N₂ⁱ) of the two NA ligands in the axial positions [symmetry code: (i) $-x + 1, -y + 1, -z + 1$] (Fig. 1).

The very similar C₁—O₁ [1.254 (2) Å] and C₁—O₂ [1.256 (2) Å], bond lengths of the carboxylate group indicate delocalized bonding arrangements, rather than localized single and double bonds. The Co—O bond lengths are 2.0835 (12) Å (for benzoate oxygen atoms) and 2.1350 (13) Å (for water oxygen atoms), and the Co—N bond length is 2.1390 (15) Å, close to standard values. The Co₁ atom lies 0.3921 (1) Å above

Table 1
Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the C₉—C₁₃,N₂ ring.

D—H···A	D—H	H···A	D···A	D—H···A
N ₃ —H ₃₁ ···O ₃ ⁱ	0.84 (3)	2.09 (3)	2.914 (3)	166 (3)
N ₃ —H ₃₂ ···O ₁ ⁱⁱ	0.87 (3)	2.13 (3)	2.910 (3)	148 (3)
O ₄ —H ₄₁ ···O ₁ ⁱⁱⁱ	0.85 (3)	1.82 (3)	2.658 (2)	166 (3)
O ₄ —H ₄₂ ···O ₃ ^{iv}	0.80 (3)	2.11 (3)	2.877 (2)	161 (2)
C ₄ —H ₄ ···O ₁ ^v	0.93	2.38	3.302 (3)	173
C ₉ —H ₉ ···N ₁ ^{vi}	0.93	2.54	3.305 (5)	140
C ₆ —H ₆ ···Cg2 ^{vii}	0.93	2.76	3.691 (2)	176

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x + 1, -y, -z + 1$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $-x, -y + 1, -z + 1$; (v) $x - 1, y, z$; (vi) $x + 1, y, z + 1$; (vii) $x, y, z - 1$.

the planar (O₁/O₂/C₁) carboxylate group. The O—Co—O and O—Co—N bond angles deviate only slightly from ideal values, with average values of 90 (3)° and 90 (2)°, respectively.


The dihedral angle between the planar carboxylate group (O₁/O₂/C₁) and the adjacent benzene ring [*A* (C₂—C₇)] is 22.11 (15)°, while the benzene and pyridine [B (N₂/C₉—C₁₃)] rings are oriented at a dihedral angle of 89.98 (5)°.

3. Supramolecular features

In the crystal, N—H···O_c (*c* = carboxylate), N—H···O_n (*n* = nicotinamide), O—H_w···O_c (*w* = water) and O—H_w···O_n hydrogen bonds (Table 1) link the molecules, enclosing R₂²(8) and R₄⁴(8) ring motifs (Bernstein *et al.*, 1995), forming layers parallel to (100) (Fig. 2). The layers are linked via C—H_{cnb}···O_c (cnb = cyanobenzoate) and C—H_n···N_{cnb} hydrogen bonds (Table 1), resulting in a three-dimensional network. A weak C—H···π interaction is also observed.

4. Synthesis and crystallization

The title compound was prepared by the reaction of CoSO₄·7H₂O (1.41 g, 5 mmol) in H₂O (50 ml) and nicotinamide (1.22 g, 50 mmol) in H₂O (50 ml) with sodium 4-cyano-

**Figure 2**

Part of the crystal structure viewed down [100], where the *b* axis is horizontal and the *c* axis is vertical. Intermolecular N—H···O and O—H···O hydrogen bonds are shown as dashed lines. Non-bonding H atoms have been omitted for clarity.

benzoate (1.69 g, 10 mmol) in H₂O (100 ml). The mixture was filtered and set aside to crystallize at ambient temperature for several days, giving pink-coloured single crystals.

5. Refinement

The experimental details including the crystal data, data collection and refinement are summarized in Table 2. Atoms H31 and H32 (for NH₂) and H41 and H42 (for H₂O) were located in a difference Fourier map and were refined freely. The aromatic C-bound H atoms were positioned geometrically with C—H = 0.93 Å, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The highest electron density and the deepest hole were found 0.80 Å and 0.83 Å, respectively, from Co1.

Acknowledgements

The authors acknowledge the Aksaray University, Science and Technology Application and Research Center, Aksaray, Turkey, for the use of the Bruker SMART BREEZE CCD diffractometer (purchased under grant No. 2010K120480 of the State Planning Organization).

References

- Adiwidjaja, G., Rossmannith, E. & Küppers, H. (1978). *Acta Cryst. B* **34**, 3079–3083.
- Amiraslanov, I. R., Mamedov, Kh. S., Movsumov, E. M., Musaev, F. N. & Nadzhafov, G. N. (1979). *Zh. Strukt. Khim.* **20**, 1075–1080.
- Antolini, L., Battaglia, L. P., Corradi, A. B., Marcotrigiano, G., Menabue, L., Pellacani, G. C. & Saladini, M. (1982). *Inorg. Chem.* **21**, 1391–1395.
- Antsyshkina, A. S., Chiragov, F. M. & Poray-Koshits, M. A. (1980). *Koord. Khim.* **15**, 1098–1103.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bigoli, F., Braibanti, A., Pellinghelli, M. A. & Tiripicchio, A. (1972). *Acta Cryst. B* **28**, 962–966.
- Bruker (2012). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc. Madison, Wisconsin, USA.
- Catterick, J., Hursthouse, M. B., New, D. B. & Thornton, P. (1974). *J. Chem. Soc. Chem. Commun.* pp. 843–844.
- Chen, H. J. & Chen, X. M. (2002). *Inorg. Chim. Acta*, **329**, 13–21.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Hauptmann, R., Kondo, M. & Kitagawa, S. (2000). *Z. Kristallogr. New Cryst. Struct.* **215**, 169–172.
- Krishnamachari, K. A. V. R. (1974). *Am. J. Clin. Nutr.* **27**, 108–111.
- Nadzhafov, G. N., Shnulin, A. N. & Mamedov, Kh. S. (1981). *Zh. Strukt. Khim.* **22**, 124–128.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shnulin, A. N., Nadzhafov, G. N., Amiraslanov, I. R., Usubaliev, B. T. & Mamedov, Kh. S. (1981). *Koord. Khim.* **7**, 1409–1416.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

Table 2
Experimental details.

Crystal data	[Co(C ₈ H ₄ NO ₂) ₂ (C ₆ H ₆ N ₂ O) ₂ (H ₂ O) ₂]
M_r	631.46
Crystal system, space group	Triclinic, $P\bar{1}$
Temperature (K)	296
a, b, c (Å)	7.6474 (3), 9.9266 (4), 10.2782 (4)
α, β, γ (°)	78.680 (2), 84.200 (3), 71.556 (2)
V (Å ³)	725.13 (5)
Z	1
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	0.65
Crystal size (mm)	0.43 × 0.29 × 0.16
Data collection	
Diffractometer	Bruker SMART BREEZE CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Bruker, 2012)
T_{\min}, T_{\max}	0.797, 0.901
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	16533, 3639, 3419
R_{int}	0.036
(sin θ/λ) _{max} (Å ⁻¹)	0.672
Refinement	
$R[F^2 > 2\sigma(F^2)]$, $wR(F^2)$, S	0.044, 0.119, 1.07
No. of reflections	3639
No. of parameters	212
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	0.94, -0.49

Computer programs: *APEX2* and *SAINT* (Bruker, 2012), *SHELXS97* and *SHELXL97* (Sheldrick, 2008), *ORTEP-3 for Windows* (Farrugia, 2012) and *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

supporting information

Acta Cryst. (2015). E71, 561-563 [doi:10.1107/S2056989015008270]

Crystal structure of *trans*-diaquabis(4-cyanobenzoato- κO)bis(nicotinamide- κN^1)cobalt(II)

Gülçin Şefiye Aşkın, Hacalı Necefoglu, Gamze Yılmaz Nayır, Raziye Çatak Çelik and Tuncer Hökelek

Computing details

Data collection: *APEX2* (Bruker, 2012); cell refinement: *SAINT* (Bruker, 2012); data reduction: *SAINT* (Bruker, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

trans-Diaquabis(4-cyanobenzoato- κO)bis(nicotinamide- κN^1)cobalt(II)

Crystal data

[Co(C ₈ H ₄ NO ₂) ₂ (C ₆ H ₆ N ₂ O) ₂ (H ₂ O) ₂]	Z = 1
M _r = 631.46	F(000) = 325
Triclinic, P $\bar{1}$	D _x = 1.446 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 7.6474 (3) Å	Cell parameters from 9898 reflections
b = 9.9266 (4) Å	θ = 2.2–28.6°
c = 10.2782 (4) Å	μ = 0.65 mm ⁻¹
α = 78.680 (2)°	T = 296 K
β = 84.200 (3)°	Prism, translucent light pink
γ = 71.556 (2)°	0.43 × 0.29 × 0.16 mm
V = 725.13 (5) Å ³	

Data collection

Bruker SMART BREEZE CCD	16533 measured reflections
diffractometer	3639 independent reflections
Radiation source: fine-focus sealed tube	3419 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.036$
φ and ω scans	$\theta_{\text{max}} = 28.6^\circ$, $\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan	$h = -10 \rightarrow 10$
(<i>SADABS</i> ; Bruker, 2012)	$k = -12 \rightarrow 13$
$T_{\text{min}} = 0.797$, $T_{\text{max}} = 0.901$	$l = -13 \rightarrow 13$

Refinement

Refinement on F^2	212 parameters
Least-squares matrix: full	0 restraints
$R[F^2 > 2\sigma(F^2)] = 0.044$	Primary atom site location: structure-invariant
$wR(F^2) = 0.119$	direct methods
$S = 1.07$	Secondary atom site location: difference Fourier
3639 reflections	map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0721P)^2 + 0.2871P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.94 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.49 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.5000	0.5000	0.5000	0.02573 (12)
O1	0.6366 (2)	0.34341 (18)	0.23257 (16)	0.0459 (4)
O2	0.38784 (18)	0.47948 (14)	0.33110 (13)	0.0347 (3)
O3	0.0668 (2)	0.16139 (15)	0.50697 (18)	0.0476 (4)
O4	0.22430 (19)	0.57799 (15)	0.57811 (16)	0.0362 (3)
H41	0.250 (4)	0.608 (3)	0.644 (3)	0.064 (9)*
H42	0.147 (4)	0.643 (3)	0.538 (2)	0.038 (6)*
N1	-0.0584 (5)	0.1958 (4)	-0.1318 (4)	0.1083 (13)
N2	0.4952 (2)	0.28815 (16)	0.59167 (15)	0.0303 (3)
N3	0.1614 (3)	-0.0568 (2)	0.6345 (2)	0.0473 (5)
H31	0.080 (4)	-0.079 (3)	0.602 (3)	0.063 (9)*
H32	0.238 (4)	-0.121 (3)	0.690 (3)	0.054 (8)*
C1	0.4662 (2)	0.40272 (19)	0.24567 (17)	0.0300 (3)
C2	0.3451 (2)	0.37166 (19)	0.15590 (17)	0.0298 (3)
C3	0.1622 (3)	0.3852 (2)	0.19215 (19)	0.0382 (4)
H3	0.1098	0.4240	0.2675	0.046*
C4	0.0562 (3)	0.3414 (3)	0.1171 (2)	0.0457 (5)
H4	-0.0661	0.3486	0.1430	0.055*
C5	0.1336 (3)	0.2869 (3)	0.0037 (2)	0.0458 (5)
C6	0.3143 (4)	0.2788 (3)	-0.0368 (2)	0.0570 (7)
H6	0.3641	0.2460	-0.1152	0.068*
C7	0.4203 (3)	0.3198 (3)	0.0398 (2)	0.0470 (5)
H7	0.5424	0.3126	0.0137	0.056*
C8	0.0255 (4)	0.2372 (4)	-0.0727 (3)	0.0680 (8)
C9	0.6293 (2)	0.1956 (2)	0.66831 (19)	0.0339 (4)
H9	0.7280	0.2252	0.6841	0.041*
C10	0.6261 (3)	0.0585 (2)	0.7244 (2)	0.0428 (5)
H10	0.7210	-0.0033	0.7772	0.051*
C11	0.4798 (3)	0.0138 (2)	0.7013 (2)	0.0419 (4)
H11	0.4756	-0.0786	0.7379	0.050*

C12	0.3394 (2)	0.10851 (18)	0.62286 (18)	0.0307 (4)
C13	0.3539 (2)	0.24444 (18)	0.57084 (18)	0.0304 (3)
H13	0.2601	0.3088	0.5185	0.037*
C14	0.1776 (3)	0.07268 (19)	0.5848 (2)	0.0352 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02515 (17)	0.02306 (18)	0.03406 (19)	-0.01019 (12)	-0.00572 (12)	-0.00978 (12)
O1	0.0315 (7)	0.0592 (9)	0.0516 (8)	-0.0103 (6)	-0.0066 (6)	-0.0234 (7)
O2	0.0355 (6)	0.0333 (6)	0.0400 (7)	-0.0084 (5)	-0.0111 (5)	-0.0157 (5)
O3	0.0445 (8)	0.0303 (7)	0.0743 (11)	-0.0156 (6)	-0.0241 (7)	-0.0061 (7)
O4	0.0290 (6)	0.0348 (7)	0.0464 (8)	-0.0075 (5)	-0.0060 (6)	-0.0120 (6)
N1	0.104 (2)	0.126 (3)	0.121 (3)	-0.030 (2)	-0.052 (2)	-0.065 (2)
N2	0.0304 (7)	0.0275 (7)	0.0370 (7)	-0.0115 (6)	-0.0065 (6)	-0.0082 (6)
N3	0.0515 (11)	0.0305 (8)	0.0691 (13)	-0.0229 (8)	-0.0187 (9)	-0.0046 (8)
C1	0.0332 (8)	0.0290 (8)	0.0312 (8)	-0.0130 (7)	-0.0068 (6)	-0.0040 (6)
C2	0.0342 (8)	0.0291 (8)	0.0288 (8)	-0.0102 (7)	-0.0060 (6)	-0.0078 (6)
C3	0.0361 (9)	0.0491 (11)	0.0349 (9)	-0.0137 (8)	-0.0023 (7)	-0.0184 (8)
C4	0.0368 (10)	0.0620 (14)	0.0475 (11)	-0.0207 (9)	-0.0054 (8)	-0.0196 (10)
C5	0.0500 (11)	0.0505 (12)	0.0437 (11)	-0.0140 (9)	-0.0159 (9)	-0.0190 (9)
C7	0.0396 (10)	0.0679 (15)	0.0403 (10)	-0.0181 (10)	0.0037 (8)	-0.0253 (10)
C6	0.0523 (13)	0.0836 (18)	0.0428 (11)	-0.0148 (12)	-0.0034 (9)	-0.0377 (12)
C8	0.0649 (16)	0.0794 (19)	0.0706 (17)	-0.0168 (14)	-0.0256 (13)	-0.0357 (15)
C9	0.0295 (8)	0.0345 (9)	0.0404 (9)	-0.0108 (7)	-0.0076 (7)	-0.0083 (7)
C10	0.0382 (10)	0.0349 (10)	0.0529 (12)	-0.0071 (8)	-0.0171 (8)	-0.0012 (8)
C11	0.0456 (11)	0.0248 (8)	0.0561 (12)	-0.0123 (8)	-0.0133 (9)	-0.0006 (8)
C12	0.0331 (8)	0.0241 (8)	0.0395 (9)	-0.0111 (6)	-0.0046 (7)	-0.0106 (7)
C13	0.0311 (8)	0.0240 (7)	0.0399 (9)	-0.0104 (6)	-0.0091 (7)	-0.0069 (7)
C14	0.0361 (9)	0.0270 (8)	0.0490 (10)	-0.0139 (7)	-0.0043 (7)	-0.0133 (7)

Geometric parameters (\AA , $^\circ$)

Co1—O2	2.0835 (12)	C2—C7	1.391 (3)
Co1—O2 ⁱ	2.0835 (12)	C3—C4	1.387 (3)
Co1—O4	2.1350 (13)	C3—H3	0.9300
Co1—O4 ⁱ	2.1350 (13)	C4—C5	1.381 (3)
Co1—N2	2.1390 (15)	C4—H4	0.9300
Co1—N2 ⁱ	2.1390 (15)	C5—C6	1.384 (4)
O1—C1	1.254 (2)	C5—C8	1.444 (3)
O2—C1	1.256 (2)	C6—H6	0.9300
O3—C14	1.234 (2)	C7—C6	1.380 (3)
O4—H41	0.85 (3)	C7—H7	0.9300
O4—H42	0.80 (3)	C9—C10	1.378 (3)
N1—C8	1.136 (4)	C9—H9	0.9300
N2—C9	1.341 (2)	C10—H10	0.9300
N2—C13	1.335 (2)	C11—C10	1.383 (3)
N3—C14	1.326 (3)	C11—H11	0.9300

N3—H31	0.84 (3)	C12—C11	1.388 (3)
N3—H32	0.87 (3)	C12—C13	1.385 (2)
C1—C2	1.506 (2)	C12—C14	1.497 (2)
C2—C3	1.381 (3)	C13—H13	0.9300
O2—Co1—O2 ⁱ	180.0	C4—C3—H3	119.7
O2—Co1—O4	87.59 (6)	C3—C4—H4	120.3
O2 ⁱ —Co1—O4	92.41 (6)	C5—C4—C3	119.5 (2)
O2—Co1—O4 ⁱ	92.41 (6)	C5—C4—H4	120.3
O2 ⁱ —Co1—O4 ⁱ	87.59 (6)	C4—C5—C6	120.46 (19)
O4—Co1—O4 ⁱ	180.000 (1)	C4—C5—C8	119.7 (2)
O2—Co1—N2	89.99 (6)	C6—C5—C8	119.8 (2)
O2 ⁱ —Co1—N2	90.01 (5)	C5—C6—H6	120.1
O2—Co1—N2 ⁱ	90.01 (5)	C7—C6—C5	119.8 (2)
O2 ⁱ —Co1—N2 ⁱ	89.99 (6)	C7—C6—H6	120.1
O4—Co1—N2	87.40 (6)	C2—C7—H7	119.9
O4 ⁱ —Co1—N2	92.60 (6)	C6—C7—C2	120.3 (2)
O4—Co1—N2 ⁱ	92.60 (6)	C6—C7—H7	119.9
O4 ⁱ —Co1—N2 ⁱ	87.40 (6)	N1—C8—C5	178.9 (4)
N2 ⁱ —Co1—N2	180.00 (4)	N2—C9—C10	122.33 (17)
C1—O2—Co1	127.68 (11)	N2—C9—H9	118.8
Co1—O4—H41	97 (2)	C10—C9—H9	118.8
Co1—O4—H42	122.7 (18)	C9—C10—C11	119.16 (17)
H42—O4—H41	107 (3)	C9—C10—H10	120.4
C9—N2—Co1	123.03 (12)	C11—C10—H10	120.4
C13—N2—Co1	118.91 (12)	C10—C11—C12	119.24 (18)
C13—N2—C9	118.05 (16)	C10—C11—H11	120.4
C14—N3—H31	116 (2)	C12—C11—H11	120.4
C14—N3—H32	123.3 (19)	C11—C12—C14	124.94 (17)
H32—N3—H31	120 (3)	C13—C12—C11	117.65 (16)
O1—C1—O2	125.70 (17)	C13—C12—C14	117.35 (16)
O1—C1—C2	116.72 (16)	N2—C13—C12	123.58 (16)
O2—C1—C2	117.48 (15)	N2—C13—H13	118.2
C3—C2—C1	120.48 (16)	C12—C13—H13	118.2
C3—C2—C7	119.43 (17)	O3—C14—N3	122.08 (18)
C7—C2—C1	119.95 (17)	O3—C14—C12	119.90 (17)
C2—C3—C4	120.52 (18)	N3—C14—C12	118.00 (18)
C2—C3—H3	119.7		
O4—Co1—O2—C1	-159.74 (16)	O2—C1—C2—C7	163.80 (19)
O4 ⁱ —Co1—O2—C1	20.26 (16)	C1—C2—C3—C4	-172.77 (19)
N2—Co1—O2—C1	-72.35 (16)	C7—C2—C3—C4	2.9 (3)
N2 ⁱ —Co1—O2—C1	107.65 (16)	C1—C2—C7—C6	174.2 (2)
O2—Co1—N2—C9	141.91 (15)	C3—C2—C7—C6	-1.6 (4)
O2 ⁱ —Co1—N2—C9	-38.09 (15)	C2—C3—C4—C5	-1.5 (4)
O2—Co1—N2—C13	-37.22 (14)	C3—C4—C5—C6	-1.4 (4)
O2 ⁱ —Co1—N2—C13	142.78 (14)	C3—C4—C5—C8	178.2 (2)
O4—Co1—N2—C9	-130.50 (15)	C4—C5—C6—C7	2.8 (4)

O4 ⁱ —Co1—N2—C9	49.50 (15)	C8—C5—C6—C7	−176.8 (3)
O4—Co1—N2—C13	50.36 (14)	C2—C7—C6—C5	−1.3 (4)
O4 ⁱ —Co1—N2—C13	−129.64 (14)	N2—C9—C10—C11	0.0 (3)
Co1—O2—C1—O1	−13.7 (3)	C12—C11—C10—C9	−0.4 (3)
Co1—O2—C1—C2	162.49 (12)	C13—C12—C11—C10	0.2 (3)
Co1—N2—C9—C10	−178.63 (15)	C14—C12—C11—C10	177.2 (2)
C13—N2—C9—C10	0.5 (3)	C11—C12—C13—N2	0.3 (3)
Co1—N2—C13—C12	178.47 (14)	C14—C12—C13—N2	−176.86 (17)
C9—N2—C13—C12	−0.7 (3)	C11—C12—C14—O3	−174.9 (2)
O1—C1—C2—C3	156.08 (19)	C11—C12—C14—N3	3.4 (3)
O1—C1—C2—C7	−19.6 (3)	C13—C12—C14—O3	2.1 (3)
O2—C1—C2—C3	−20.5 (3)	C13—C12—C14—N3	−179.63 (19)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , °)

Cg2 is the centroid of the C9—C13,N2 ring.

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N3—H31…O3 ⁱⁱ	0.84 (3)	2.09 (3)	2.914 (3)	166 (3)
N3—H32…O1 ⁱⁱⁱ	0.87 (3)	2.13 (3)	2.910 (3)	148 (3)
O4—H41…O1 ⁱ	0.85 (3)	1.82 (3)	2.658 (2)	166 (3)
O4—H42…O3 ^{iv}	0.80 (3)	2.11 (3)	2.877 (2)	161 (2)
C4—H4…O1 ^v	0.93	2.38	3.302 (3)	173
C9—H9…N1 ^{vi}	0.93	2.54	3.305 (5)	140
C6—H6…Cg2 ^{vii}	0.93	2.76	3.691 (2)	176

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y, -z+1$; (iii) $-x+1, -y, -z+1$; (iv) $-x, -y+1, -z+1$; (v) $x-1, y, z$; (vi) $x+1, y, z+1$; (vii) $x, y, z-1$.