

ON GENERALIZED CLASS OF p -VALENT FUNCTIONS WITH NEGATIVE COEFFICIENTS

R.M. EL-ASHWAH¹, M.K. AOUF², H.M. ZAYED³

ABSTRACT. In this paper we introduce and study new class $F_{p,\theta}(\gamma, \beta)$ of p -valent functions with negative coefficients. We obtain coefficients inequalities, distortion theorems, extreme points and radii of close to convexity, starlikeness and convexity for the class $F_{p,\theta}(\gamma, \beta)$. Also modified Hadamard products of several functions belonging to the class $F_{p,\theta}(\gamma, \beta)$ are study here. Finally, we investigate several distortion inequalities involving fractional calculus.

Keywords: Analytic, p -valent functions, Hadamard product, fractional calculus operators.

AMS Subject Classification: 30C45.

1. INTRODUCTION

Let $T_p(\theta)$ denote the class of functions of the form:

$$f(z) = z^p - \sum_{k=1}^{\infty} a_{p+k} z^{p+k} \quad (e^{i\theta} a_{p+k} \geq 0; |\theta| < \frac{\pi}{2}; p \in \mathbb{N} = \{1, 2, \dots\}), \quad (1)$$

which are analytic and p -valent in the open unit disc $U = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$. Also, let $F_{p,\theta}(\gamma, \beta)$ denote the class of functions $f(z) \in T_p(\theta)$ which satisfy

$$\operatorname{Re} \left\{ e^{i\theta} \left((1 - \gamma) \frac{f(z)}{z^p} + \gamma \frac{f'(z)}{pz^{p-1}} \right) \right\} > \frac{\beta}{p}, \quad (2)$$

where $0 \leq \frac{\beta}{p} < \cos \theta$, $|\theta| < \frac{\pi}{2}$, $\gamma \geq 0$, $p \in \mathbb{N}$ and $z \in U$.

We note that for suitable choices of γ, θ and p , we obtain the following subclasses:

- (1) $F_{p,0}(\gamma, \beta) = F_p(\gamma, \beta)$ ($0 \leq \beta < p$, $\gamma \geq 0$, $p \in \mathbb{N}$) (see Lee et al. [4] and Aouf and Darwish [2]);
- (2) $F_{1,0}(\gamma, \beta) = F(\gamma, \beta)$ ($0 \leq \beta < 1$, $\gamma \geq 0$) (see Bhoosnurmath and Swamy [3]);
- (3) $F_{1,\theta}(1, \beta) = A(\theta, \beta)$ ($0 \leq \beta < \cos \theta$, $|\theta| < \frac{\pi}{2}$) (see Sekine [7]).

2. COEFFICIENT ESTIMATES

Unless otherwise mentioned, we assume throughout this paper that

$$e^{i\theta} a_{p+k} \geq 0, \quad 0 \leq \frac{\beta}{p} < \cos \theta, \quad |\theta| < \frac{\pi}{2}, \quad \gamma \geq 0 \text{ and } p, k \in \mathbb{N}.$$

¹Department of Mathematics, Faculty of Science, Damietta University, New Damietta 34517, Egypt,

²Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt,

³Department of Mathematics, Faculty of Science, Menofia University, Shebin Elkom 32511, Egypt,

e-mail: r.elashwah@yahoo.com, mkaouf127@yahoo.com, hanaazayed42@yahoo.com

Manuscript received February 2013.

Theorem 2.1. *Let the function $f(z)$ be given by (1.1). Then $f(z) \in F_{p,\theta}(\gamma, \beta)$ if and only if*

$$\sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k} \leq \cos \theta - \frac{\beta}{p}. \tag{3}$$

Proof. Assume that the condition (3) holds true, then it is sufficient to show that the value for

$$e^{i\theta} \left((1 - \gamma) \frac{f(z)}{z^p} + \gamma \frac{f'(z)}{pz^{p-1}} \right),$$

lie in a circle centered at a point $e^{i\theta}$ whose radius is $\cos \theta - \frac{\beta}{p}$. Indeed, we have

$$\begin{aligned} & \left| e^{i\theta} \left((1 - \gamma) \frac{f(z)}{z^p} + \gamma \frac{f'(z)}{pz^{p-1}} \right) - e^{i\theta} \right| = \\ & = \left| e^{i\theta} \sum_{k=1}^{\infty} \left(1 + \frac{\gamma k}{p}\right) a_{p+k} z^k \right| \leq \\ & \leq \sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k} \leq \\ & \leq \cos \theta - \frac{\beta}{p}. \end{aligned}$$

Conversely, assume that

$$\operatorname{Re} \left\{ e^{i\theta} \left((1 - \gamma) \frac{f(z)}{z^p} + \gamma \frac{f'(z)}{pz^{p-1}} \right) \right\} > \frac{\beta}{p},$$

which is equivalent to

$$\operatorname{Re} \left\{ \sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k} z^k \right\} < \cos \theta - \frac{\beta}{p}.$$

Choose values of z on the real axis so that

$$\sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k} z^k,$$

is real. Letting $z \rightarrow 1^-$ along the real axis, we have

$$\sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k} \leq \cos \theta - \frac{\beta}{p},$$

and hence the proof of Theorem 2.1 is completed. □

Corollary 2.1. *Let the function $f(z)$ defined by (1.1) be in the class $F_{p,\theta}(\gamma, \beta)$. Then*

$$|a_{p+k}| \leq \frac{p \cos \theta - \beta}{p + k\gamma}. \tag{4}$$

The result is sharp for the function

$$f(z) = z^p - \frac{p \cos \theta - \beta}{p + k\gamma} e^{-i\theta} z^{p+k}. \tag{5}$$

3. DISTORTION THEOREMS

Theorem 3.1. *Let the function $f(z)$ defined by (1) be in the class $F_{p,\theta}(\gamma, \beta)$, then for $z \in U$, we have*

$$|z|^p - \frac{p \cos \theta - \beta}{p + \gamma} |z|^{p+1} \leq |f(z)| \leq |z|^p + \frac{p \cos \theta - \beta}{p + \gamma} |z|^{p+1}. \quad (6)$$

Furthermore

$$p |z|^{p-1} - \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)} |z|^p \leq |f'(z)| \leq p |z|^{p-1} + \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)} |z|^p. \quad (7)$$

The result is sharp for the function $f(z)$ given by

$$f(z) = z^p - \frac{p \cos \theta - \beta}{p + \gamma} e^{-i\theta} z^{p+1} \quad (z = \pm |z| e^{i\theta}). \quad (8)$$

Proof. It is easy to see from Theorem 2.1 that

$$\frac{p + \gamma}{p \cos \theta - \beta} \sum_{k=1}^{\infty} |a_{p+k}| \leq \sum_{k=1}^{\infty} \frac{p + k\gamma}{p \cos \theta - \beta} |a_{p+k}| \leq 1.$$

Then

$$\sum_{k=1}^{\infty} |a_{p+k}| \leq \frac{p \cos \theta - \beta}{p + \gamma}. \quad (9)$$

Making use of (9), we have

$$|f(z)| \geq |z|^p - |z|^{p+1} \sum_{k=1}^{\infty} |a_{p+k}| \geq |z|^p - \frac{p \cos \theta - \beta}{p + \gamma} |z|^{p+1}, \quad (10)$$

and

$$|f(z)| \leq |z|^p + |z|^{p+1} \sum_{k=1}^{\infty} |a_{p+k}| \leq |z|^p + \frac{p \cos \theta - \beta}{p + \gamma} |z|^{p+1}, \quad (11)$$

which proves the assertion (6).

From (9) and Theorem 2.1, it follows also that

$$\sum_{k=1}^{\infty} (p+k) |a_{p+k}| \leq \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)}. \quad (12)$$

Consequently, we have

$$|f'(z)| \geq p |z|^{p-1} - |z|^p \sum_{k=1}^{\infty} (p+k) |a_{p+k}| \geq p |z|^{p-1} - \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)} |z|^p, \quad (13)$$

and

$$|f'(z)| \leq p |z|^{p-1} + |z|^p \sum_{k=1}^{\infty} (p+k) |a_{p+k}| \leq p |z|^{p-1} + \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)} |z|^p, \quad (14)$$

which proves the assertion (7). Since each of equalities in (6) and (7) is satisfied by the function $f(z)$ given by (8), our proof of Theorem 3.1 is thus completed. \square

4. CLOSURE THEOREMS

Let the functions $f_j(z)$ be defined, for $j = 1, 2, \dots, m$, by

$$f_j(z) = z^p - \sum_{k=1}^{\infty} a_{p+k,j} z^{p+k} \quad (e^{i\theta} a_{p+k,j} \geq 0; |\theta| < \frac{\pi}{2}). \tag{15}$$

Theorem 4.1. *Let the functions $f_j(z)$ ($j = 1, 2, \dots, m$) defined by (15) be in the class $F_{p,\theta}(\gamma, \beta)$. Then the function $h(z)$ defined by*

$$h(z) = z^p - \sum_{k=1}^{\infty} b_{p+k} z^{p+k}, \tag{16}$$

also belongs to the class $F_{p,\theta}(\gamma, \beta)$, where

$$b_{p+k} = \frac{1}{m} \sum_{j=1}^m a_{p+k,j}. \tag{17}$$

Proof. Since $f_j(z)$ ($j = 1, 2, \dots, m$) are in the class $F_{p,\theta}(\gamma, \beta)$, it follows from Theorem 2.1, that

$$\sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k,j} \leq \cos \theta - \frac{\beta}{p},$$

for every $j = 1, 2, \dots, m$. Hence

$$\begin{aligned} \sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) b_{p+k} &= \sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) \left(\frac{1}{m} \sum_{j=1}^m a_{p+k,j}\right) = \\ &= \frac{1}{m} \sum_{j=1}^m \left(\sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k,j}\right) \leq \frac{1}{m} \sum_{j=1}^m \left(\cos \theta - \frac{\beta}{p}\right) \leq \cos \theta - \frac{\beta}{p}. \end{aligned}$$

By Theorem 2.1, it follows that $h(z) \in F_{p,\theta}(\gamma, \beta)$. This completes the proof of Theorem 4.1. \square

Theorem 4.2. *Let the functions $f_j(z)$ ($j = 1, 2, \dots, m$) defined by (15) be in the class $F_{p,\theta}(\gamma, \beta_j)$. Then the function $h(z)$ defined by*

$$h(z) = z^p - \sum_{k=1}^{\infty} \left(\frac{1}{m} \sum_{j=1}^m a_{p+k,j}\right) z^{p+k}, \tag{18}$$

is in the class $F_{p,\theta}(\gamma, \beta)$, where

$$\beta = \min_{1 \leq j \leq m} \{\beta_j\}. \tag{19}$$

Proof. Since $f_j(z)$ ($j = 1, 2, \dots, m$) are in the class $F_{p,\theta}(\gamma, \beta_j)$, it follows from Theorem 2.1, that

$$\sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k,j} \leq \cos \theta - \frac{\beta_j}{p},$$

for every $j = 1, 2, \dots, m$. Hence

$$\begin{aligned} \sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) \left(\frac{1}{m} \sum_{j=1}^m a_{p+k,j}\right) &= \frac{1}{m} \sum_{j=1}^m \left(\sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p}\right) a_{p+k,j}\right) \leq \\ &\leq \frac{1}{m} \sum_{j=1}^m \left(\cos \theta - \frac{\beta_j}{p}\right) \leq \frac{1}{m} \sum_{j=1}^m \left(\cos \theta - \frac{\beta}{p}\right) \leq \cos \theta - \frac{\beta}{p}. \end{aligned}$$

By Theorem 2.1, it follows that $h(z) \in F_{p,\theta}(\gamma, \beta)$. This completes the proof of Theorem 4.2. \square

Theorem 4.3. Let the functions $f_j(z)$ ($j = 1, 2, \dots, m$) defined by (15) be in the class $F_{p,\theta}(\gamma, \beta)$. Then the function $h(z)$ defined by

$$h(z) = \sum_{j=1}^m c_j f_j(z), \quad (20)$$

is also in the class $F_{p,\theta}(\gamma, \beta)$, where

$$\sum_{j=1}^m c_j = 1. \quad (21)$$

Proof. Assume that

$$h(z) = \sum_{j=1}^m c_j f_j(z) = z^p - \sum_{k=1}^{\infty} \left(\sum_{j=1}^m c_j a_{p+k,j} \right) z^{p+k}. \quad (22)$$

Then it follows that

$$\begin{aligned} \sum_{k=1}^{\infty} \left(1 + \frac{\gamma k}{p} \right) e^{i\theta} \left(\sum_{j=1}^m c_j a_{p+k,j} \right) &= \sum_{j=1}^m c_j \left(\sum_{k=1}^{\infty} e^{i\theta} \left[1 + \frac{\gamma k}{p} \right] a_{p+k,j} \right) \leq \\ &\leq \left(\cos \theta - \frac{\beta}{p} \right) \sum_{j=1}^m c_j \leq \cos \theta - \frac{\beta}{p}. \end{aligned}$$

By Theorem 2.1, it follows that $h(z) \in F_{p,\theta}(\gamma, \beta)$. This completes the proof of Theorem 4.3. \square

Theorem 4.4. Let $f_p(z) = z^p$ and

$$f_{p+k}(z) = z^p - \frac{p \cos \theta - \beta}{p + k\gamma} e^{-i\theta} z^{p+k}. \quad (23)$$

Then $f(z)$ is in the class $F_{p,\theta}(\gamma, \beta)$ if and only if can be expressed in the form

$$f(z) = \sum_{k=0}^{\infty} \mu_{p+k} f_{p+k}(z), \quad (24)$$

where $\mu_{p+k} \geq 0$ and $\sum_{k=0}^{\infty} \mu_{p+k} = 1$.

Proof. Assume that

$$f(z) = \sum_{k=0}^{\infty} \mu_{p+k} f_{p+k}(z) = z^p - \sum_{k=1}^{\infty} \frac{p \cos \theta - \beta}{p + k\gamma} e^{-i\theta} \mu_{p+k} z^{p+k}. \quad (25)$$

Then it follows that

$$\begin{aligned} \sum_{k=1}^{\infty} e^{i\theta} \left(1 + \frac{\gamma k}{p} \right) \left(\frac{p \cos \theta - \beta}{p + k\gamma} \right) e^{-i\theta} \mu_{p+k} &= \left(\cos \theta - \frac{\beta}{p} \right) \sum_{k=1}^{\infty} \mu_{p+k} = \\ &= \left(\cos \theta - \frac{\beta}{p} \right) (1 - \mu_p) \leq \cos \theta - \frac{\beta}{p}, \end{aligned}$$

which implies that $f(z) \in F_{p,\theta}(\gamma, \beta)$.

Conversely, assume that the function $f(z)$ defined by (1) be in the class $F_{p,\theta}(\gamma, \beta)$. Then

$$a_{p+k} \leq \frac{(p \cos \theta - \beta) e^{-i\theta}}{(p + k\gamma)}.$$

Setting

$$\mu_{p+k} = \frac{(p + k\gamma)}{(p \cos \theta - \beta)} e^{i\theta} a_{p+k},$$

where

$$\mu_p = 1 - \sum_{k=1}^{\infty} \mu_{p+k} ,$$

we can see that $f(z)$ can be expressed in the form (24). This completes the proof of Theorem 4.4. □

Corollary 4.1. *The extreme points of the class $F_{p,\theta}(\gamma, \beta)$ are the functions $f_p(z) = z^p$ and*

$$f_{p+k}(z) = z^p - \frac{p \cos \theta - \beta}{p + k\gamma} e^{-i\theta} z^{p+k} . \tag{26}$$

5. RADII OF CLOSE-TO-CONVEXITY, STARLIKENESS AND CONVEXITY

Theorem 5.1. *Let the function $f(z)$ defined by (1) be in the class $F_{p,\theta}(\gamma, \beta)$. Then $f(z)$ is p -valent close-to-convex of order δ ($0 \leq \delta < p$) in $|z| \leq r_1$, where*

$$r_1 = \inf_{k \geq 1} \left\{ \frac{(p + k\gamma)(p - \delta)}{(p \cos \theta - \beta)(k + p)} \right\}^{\frac{1}{k}} . \tag{27}$$

The result is sharp and the extremal function is given by (5).

Proof. We must show that

$$\left| \frac{f'(z)}{z^{p-1}} - p \right| \leq p - \delta \text{ for } |z| \leq r_1, \tag{28}$$

where r_1 is given by (27). Indeed we find from (1) that

$$\left| \frac{f'(z)}{z^{p-1}} - p \right| \leq \sum_{k=1}^{\infty} (p + k) |a_{p+k}| |z|^k .$$

Thus

$$\left| \frac{f'(z)}{z^{p-1}} - p \right| \leq p - \delta,$$

if

$$\sum_{k=1}^{\infty} \left(\frac{k + p}{p - \delta} \right) |a_{p+k}| |z|^k \leq 1. \tag{29}$$

But by using Theorem 2.1, (29) will be true if

$$\left(\frac{k + p}{p - \delta} \right) |z|^k \leq \left(\frac{p + k\gamma}{p \cos \theta - \beta} \right) .$$

Then

$$|z| \leq \left\{ \frac{(p + k\gamma)(p - \delta)}{(p \cos \theta - \beta)(k + p)} \right\}^{\frac{1}{k}} . \tag{30}$$

The result follows easily from (30). □

Theorem 5.2. Let the function $f(z)$ defined by (1) be in the class $F_{p,\theta}(\gamma, \beta)$. Then $f(z)$ is p -valent starlike of order δ ($0 \leq \delta < p$) in $|z| \leq r_2$, where

$$r_2 = \inf_{k \geq 1} \left\{ \frac{(p + k\gamma)(p - \delta)}{(p \cos \theta - \beta)(k + p - \delta)} \right\}^{\frac{1}{k}}. \quad (31)$$

The result is sharp and the extremal function is given by (5).

Proof. We must show that

$$\left| \frac{zf'(z)}{f(z)} - p \right| \leq p - \delta \text{ for } |z| \leq r_2, \quad (32)$$

where r_2 is given by (31). Indeed we find from (1) that

$$\left| \frac{zf'(z)}{f(z)} - p \right| \leq \frac{\sum_{k=1}^{\infty} k |a_{p+k}| |z|^k}{1 - \sum_{k=1}^{\infty} |a_{p+k}| |z|^k}.$$

Thus

$$\left| \frac{zf'(z)}{f(z)} - p \right| \leq p - \delta,$$

if

$$\sum_{k=1}^{\infty} \left(\frac{k + p - \delta}{p - \delta} \right) |a_{p+k}| |z|^k \leq 1. \quad (33)$$

But by using Theorem 2.1, (33) will be true if

$$\left(\frac{k + p - \delta}{p - \delta} \right) |z|^k \leq \left(\frac{p + k\gamma}{p \cos \theta - \beta} \right).$$

Then

$$|z| \leq \left\{ \frac{(p + k\gamma)(p - \delta)}{(p \cos \theta - \beta)(k + p - \delta)} \right\}^{\frac{1}{k}}. \quad (34)$$

The result follows easily from (34). \square

Corollary 5.1. Let the function $f(z)$ defined by (1) be in the class $F_{p,\theta}(\gamma, \beta)$. Then $f(z)$ is in p -valent convex of order δ ($0 \leq \delta < p$) in $|z| \leq r_3$, where

$$r_3 = \inf_{k \geq 1} \left\{ \frac{p(p + k\gamma)(p - \delta)}{(k + p)(p \cos \theta - \beta)(k + p - \delta)} \right\}^{\frac{1}{k}}. \quad (35)$$

The result is sharp and the extremal function is given by (5).

6. MODIFIED HADAMARD PRODUCTS

For the functions $f_j(z)$ ($j = 1, 2$) defined by (15) and belonging to the class $T_p(\theta)$, the modified Hadamard product of $f_1(z)$ and $f_2(z)$ is defined by

$$(f_1 * f_2)(z) = z^p - \sum_{k=1}^{\infty} a_{p+k,1} a_{p+k,2} z^{p+k}. \quad (36)$$

Theorem 6.1. *Let the functions $f_j(z)$ ($j = 1, 2$) defined by (15) be in the class $F_{p,\theta}(\gamma, \beta)$. Then $(f_1 * f_2)(z) \in F_{p,2\theta}(\gamma, \alpha)$ where*

$$\alpha = p \cos 2\theta - \frac{(p \cos \theta - \beta)^2}{(p + \gamma)}. \tag{37}$$

The result is sharp for the functions $f_j(z)$ given by

$$f_j(z) = z^p - \left(\frac{p \cos \theta - \beta}{p + \gamma} \right) e^{-i\theta} z^{p+1} \quad (j = 1, 2). \tag{38}$$

Proof. Employing the technique used earlier by Schild and Silverman [6]. We need only to find the largest α such that

$$\sum_{k=1}^{\infty} e^{2i\theta} \left(\frac{p + k\gamma}{p \cos 2\theta - \alpha} \right) a_{p+k,1} a_{p+k,2} \leq 1. \tag{39}$$

Since $f_j(z)$ ($j = 1, 2$) are in the class $F_{p,\theta}(\gamma, \beta)$, it follows from Theorem 2.1, that

$$\sum_{k=1}^{\infty} e^{i\theta} \left(\frac{p + k\gamma}{p \cos \theta - \beta} \right) a_{p+k,j} \leq 1, \tag{40}$$

for every $j = 1, 2$. By the Cauchy Schwarz inequality we have

$$\sum_{k=1}^{\infty} e^{i\theta} \left(\frac{p + k\gamma}{p \cos \theta - \beta} \right) \sqrt{a_{p+k,1} a_{p+k,2}} \leq 1. \tag{41}$$

Therefore, (39) will be satisfied if

$$e^{2i\theta} \left(\frac{p + k\gamma}{p \cos 2\theta - \alpha} \right) a_{p+k,1} a_{p+k,2} \leq e^{i\theta} \left(\frac{p + k\gamma}{p \cos \theta - \beta} \right) \sqrt{a_{p+k,1} a_{p+k,2}}.$$

Then

$$\sqrt{a_{p+k,1} a_{p+k,2}} \leq \left(\frac{p \cos 2\theta - \alpha}{p \cos \theta - \beta} \right) e^{-i\theta}. \tag{42}$$

Since (41) implies

$$\sqrt{a_{p+k,1} a_{p+k,2}} \leq \left(\frac{p \cos \theta - \beta}{p + k\gamma} \right) e^{-i\theta}. \tag{43}$$

From ((42) and (43) we have

$$\alpha \leq p \cos 2\theta - \frac{(p \cos \theta - \beta)^2}{(p + k\gamma)}. \tag{44}$$

Now defining the function $G(k)$ by

$$G(k) = p \cos 2\theta - \frac{(p \cos \theta - \beta)^2}{(p + k\gamma)}, \tag{45}$$

we see that $G(k)$ is an increasing function of k ($k \in \mathbb{N}$). Therefore, we conclude that

$$\alpha \leq G(1) = p \cos 2\theta - \frac{(p \cos \theta - \beta)^2}{(p + \gamma)}, \tag{46}$$

which evidently completes the proof of Theorem 6.1. □

Using arguments similar to those in the proof of Theorem 6.1, we obtain the following theorem.

Theorem 6.2. Let the function $f_1(z)$ defined by (15) be in the class $F_{p,\theta}(\gamma, \beta)$. Suppose also that the function $f_2(z)$ defined by (15) be in the class $F_{p,\theta}(\gamma, \phi)$. Then $(f_1 * f_2)(z) \in F_{p,2\theta}(\gamma, \zeta)$, where

$$\zeta = p \cos 2\theta - \frac{(p \cos \theta - \beta)(p \cos \theta - \phi)}{(p + \gamma)}. \quad (47)$$

The result is sharp for the functions $f_j(z)$ ($j = 1, 2$) given by

$$f_1(z) = z^p - \left(\frac{p \cos \theta - \beta}{p + \gamma} \right) e^{-i\theta} z^{p+1}, \quad (48)$$

and

$$f_2(z) = z^p - \left(\frac{p \cos \theta - \phi}{p + \gamma} \right) e^{-i\theta} z^{p+1}. \quad (49)$$

Theorem 6.3. . Let the functions $f_j(z)$ ($j = 1, 2$) defined by (15) be in the class $F_{p,\theta}(\gamma, \beta)$. Then the function

$$h(z) = z^p - \sum_{k=1}^{\infty} (a_{p+k,1}^2 + a_{p+k,2}^2) z^{p+k}, \quad (50)$$

also belongs to the class $F_{p,2\theta}(\gamma, \eta)$, where

$$\eta(p; \beta, \gamma; \theta) = p \cos 2\theta - \frac{2(p \cos \theta - \beta)^2}{(p + \gamma)}. \quad (51)$$

The result is sharp for the functions given by (38).

Proof. By using Theorem 2.1, we have

$$\sum_{k=1}^{\infty} \left[e^{i\theta} \frac{p + k\gamma}{p \cos \theta - \beta} \right]^2 a_{p+k,1}^2 \leq \left[\sum_{k=1}^{\infty} e^{i\theta} \frac{p + k\gamma}{p \cos \theta - \beta} a_{p+k,1} \right]^2 \leq 1, \quad (52)$$

and

$$\sum_{k=1}^{\infty} \left[e^{i\theta} \frac{p + k\gamma}{p \cos \theta - \beta} \right]^2 a_{p+k,2}^2 \leq \left[\sum_{k=1}^{\infty} e^{i\theta} \frac{p + k\gamma}{p \cos \theta - \beta} a_{p+k,2} \right]^2 \leq 1. \quad (53)$$

It follow from (52) and (53) that

$$\sum_{k=1}^{\infty} \frac{1}{2} \left[e^{i\theta} \frac{p + k\gamma}{p \cos \theta - \beta} \right]^2 (a_{p+k,1}^2 + a_{p+k,2}^2) \leq 1. \quad (54)$$

Therefore, we need to find the largest η such that

$$e^{2i\theta} \frac{p + k\gamma}{p \cos 2\theta - \eta} \leq \frac{1}{2} \left[e^{i\theta} \frac{p + k\gamma}{p \cos \theta - \beta} \right]^2, \quad (55)$$

that is

$$\eta \leq p \cos 2\theta - \frac{2(p \cos \theta - \beta)^2}{(p + k\gamma)}.$$

Since

$$D(k) = p \cos 2\theta - \frac{2(p \cos \theta - \beta)^2}{(p + k\gamma)},$$

is an increasing function of k ($k \in \mathbb{N}$), we obtain

$$\eta \leq D(1) = p \cos 2\theta - \frac{2(p \cos \theta - \beta)^2}{(p + \gamma)},$$

and hence the proof of Theorem 6.3 is completed. \square

Remark 6.1. (1) Putting $\theta = 0$ in our results, we obtain the results obtained by Lee et al. [4];
 (2) Putting $\gamma = p = 1$ in our results, we obtain the results obtained by Sekine [7].

7. DEFINITIONS AND APPLICATIONS OF FRACTIONAL CALCULUS

Many essentially equivalent definitions of fractional calculus (that is, fractional derivatives and fractional integrals) have been given in the literature (cf., e.g. [1], [9] and [10]. We find it to be convenient to recall here the following definitions which were used recently by Owa [5] and by Srivastava and Owa [8]).

Definition 7.1. The fractional integral of order μ is defined, for a function $f(z)$, by

$$D_z^{-\mu} f(z) = \frac{1}{\Gamma(\mu)} \int_0^z \frac{f(t)}{(z-t)^{1-\mu}} dt \quad (\mu > 0), \tag{56}$$

where $f(z)$ is an analytic function in a simply-connected region of the complex z -plane containing the origin and the multiplicity of $(z-t)^{\mu-1}$ is removed by requiring $\log(z-t)$ to be real when $z-t > 0$.

Definition 7.2. The fractional derivative of order μ is defined, for a function $f(z)$, by

$$D_z^\mu f(z) = \frac{1}{\Gamma(1-\mu)} \frac{d}{dz} \int_0^z \frac{f(t)}{(z-t)^\mu} dt \quad (0 \leq \mu < 1), \tag{57}$$

where $f(z)$ is an analytic function in a simply-connected region of the complex z -plane containing the origin and the multiplicity of $(z-t)^{-\mu}$ is removed by requiring $\log(z-t)$ to be real when $z-t > 0$.

Definition 7.3. Under the hypotheses of definition 2, the fractional derivative of order $n + \mu$ is defined by

$$D_z^{n+\mu} f(z) = \frac{d^n}{dz^n} D_z^\mu f(z) \quad (0 \leq \mu < 1; n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}). \tag{58}$$

Theorem 7.1. Let the function $f(z)$ defined by (1) be in the class $F_{p,\theta}(\gamma, \beta)$. Then we have

$$|D_z^{-\mu} f(z)| \geq \frac{\Gamma(p+1)}{\Gamma(p+\mu+1)} |z|^{p+\mu} \left\{ 1 - \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)(p+\mu+1)} |z| \right\}, \tag{59}$$

and

$$|D_z^{-\mu} f(z)| \leq \frac{\Gamma(p+1)}{\Gamma(p+\mu+1)} |z|^{p+\mu} \left\{ 1 + \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)(p+\mu+1)} |z| \right\}, \tag{60}$$

for $\mu > 0$ and $z \in U$. The result is sharp.

Proof. Let

$$\begin{aligned} F(z) &= \frac{\Gamma(p+\mu+1)}{\Gamma(p+1)} z^{-\mu} D_z^{-\mu} f(z) \\ &= z^p - \sum_{k=1}^{\infty} \frac{\Gamma(p+k+1)\Gamma(p+\mu+1)}{\Gamma(p+1)\Gamma(p+k+\mu+1)} a_{p+k} z^{p+k}. \end{aligned}$$

Then

$$F(z) = z^p - \sum_{k=1}^{\infty} \Psi(k) a_{p+k} z^{p+k}, \tag{61}$$

where

$$\Psi(k) = \frac{\Gamma(p+k+1)\Gamma(p+\mu+1)}{\Gamma(p+1)\Gamma(p+k+\mu+1)} \quad (\mu > 0).$$

Since $\Psi(k)$ is an decreasing function of k ($k \in \mathbb{N}$), then

$$0 < \Psi(k) \leq \Psi(1) = \frac{(p+1)}{(p+\mu+1)}. \quad (62)$$

From (61) and (62), we have

$$|F(z)| \geq |z|^p - \Psi(1) |z|^{p+1} \sum_{k=1}^{\infty} |a_{p+k}|. \quad (63)$$

In view of (9) and (63), we have

$$|F(z)| = \left| \frac{\Gamma(p+\mu+1)}{\Gamma(p+1)} z^{-\mu} D_z^{-\mu} f(z) \right| \geq |z|^p - \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)(p+\mu+1)} |z|^{p+1},$$

and

$$|F(z)| = \left| \frac{\Gamma(p+\mu+1)}{\Gamma(p+1)} z^{-\mu} D_z^{-\mu} f(z) \right| \leq |z|^p + \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)(p+\mu+1)} |z|^{p+1}.$$

which proves the inequalities of Theorem 7.1. Further equalities are attained for the function

$$D_z^{-\mu} f(z) = \frac{\Gamma(p+1)}{\Gamma(p+\mu+1)} z^{p+\mu} \left\{ 1 - \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)(p+\mu+1)} z \right\}, \quad (64)$$

or

$$f(z) = z^p - \frac{p \cos \theta - \beta}{p+\gamma} e^{-i\theta} z^{p+1} (z = \pm |z| e^{i\theta}). \quad (65)$$

□

Using arguments similar to those in the proof of Theorem 7.1, we obtain the following theorem.

Theorem 7.2. *Let the function $f(z)$ defined by (1) be in the class $F_{p,\theta}(\gamma, \beta)$. Then we have*

$$|D_z^{\mu} f(z)| \geq \frac{\Gamma(p+1)}{\Gamma(p-\mu+1)} |z|^{p-\mu} \left\{ 1 - \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)(p-\mu+1)} |z| \right\}, \quad (66)$$

and

$$|D_z^{\mu} f(z)| \leq \frac{\Gamma(p+1)}{\Gamma(p-\mu+1)} |z|^{p-\mu} \left\{ 1 + \frac{(p+1)(p \cos \theta - \beta)}{(p+\gamma)(p-\mu+1)} |z| \right\}, \quad (67)$$

for $0 \leq \mu < 1$ and $z \in U$. The result is sharp for the function $f(z)$ given by (65).

Remark 7.1. (1) Putting $\theta = 0$ in our results, we obtain the results obtained by Aouf and Darwish [2];

(2) Putting $\theta = 0$ and $p = 1$ in our results, we obtain the results obtained by Bhoosnurmath and Swamy [3].

REFERENCES

- [1] Aouf, M. K., (1990), On fractional derivatives and fractional integrals of certain subclasses of starlike and convex functions, Math. Japon. 35, no. 5, pp. 831-837.
- [2] Aouf, M. K., Darwish, H. E., (1994), Basic properties and characterizations of a certain class of analytic functions with negative coefficients . II, Utilitas Math., 46, pp.167-177.
- [3] Bhoosnurmath, S. S., Swamy, S. R., (1985), Certain classes of analytic functions with negative coefficients, Indian J. Math. 27, pp.89-98.
- [4] Lee, S. K., Owa, S., Srivastava, H. M., (1989), Basic properties and characterizations of a certain class of analytic functions with negative coefficients, Utilitas Math., 36, pp.121-128.
- [5] Owa, S., (1978), On the distortion theorems. I, Kyungpook Math. J. 18, pp.53-59.
- [6] Schild, A., Silverman, H., (1975), Convolutions of univalent functions with negative coefficients, Ann. Univ. Mariae-Curie Sklodowska Sect. A, 29, pp.109-116.

- [7] Sekine, T., (1991), On generalized class of analytic functions with negative coefficients, Math. Japon. 36, 1, pp.13-19.
 - [8] Srivastava, H. M., Owa, S., (1984), An application of the fractional derivative, Math. Japon. 29, pp.383-389.
 - [9] Srivastava, H. M., Owa, S., (Editors), (1989), Univalent Functions, Fractional Calculus, and Their Applications, Halsted press (Ellis Horwood Limited Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto.
 - [10] Srivastava, H. M., Aouf, M. K., (1995), A certain fractional derivative operator and its applications to a new class of analytic and multivalent functions with negative coefficients. I and II, J. Math. Anal. Appl., 171(1992), 1-13; ibid. 192, pp.973-688.
-
-

Rabha Mohammed El-Ashwah, for a photograph and biography, see TWMS J. Pure Appl. Math., V.4, N.1, 2013, p.101.

Mohammed Kamal Aouf, for a photograph and biography, see TWMS J. Pure Appl. Math., V.4, N.1, 2013, p.101.

Hanaa Mousa Ahmed Zayed was born in 1986, in Kafr El-Sheikh (Egypt). She works as assistant lecturer, Department of Mathematics, Faculty of Science, Menofia University, Egypt. Her research interest is Complex Analysis and Functional Analysis.