

## ON A HYBRID NUMERICAL ALGORITHM FOR THE SOLUTIONS OF HIGHER ORDER ORDINARY DIFFERENTIAL EQUATIONS

TIMOTHY A. ANAKE<sup>1</sup>, SHEILA A. BISHOP<sup>1</sup>, OLASUNMBO O. AGBOOLA<sup>1</sup>

**ABSTRACT.** A hybrid Adam-Moulton type one step numerical algorithm is proposed in this paper. The numerical algorithm is implemented in the block mode. Characterization of the method in terms of convergence and region of stability is given. Numerical experiments performed reveals the convergence of the method at very reasonable cost.

**Keywords:** convergence, hybrid, block method, algorithm, stability.

**AMS Subject Classification:** 65D30, 65L05, 65L06, 65L07.

### 1. INTRODUCTION

We propose a numerical algorithm for the solution of higher order ordinary differential equations(ODE), of the form:

$$\begin{cases} \ddot{y} = f(x, y(x), \dot{y}(x)), & x \in [a, T] \\ y(a) = \xi_0, \\ \dot{y}(a) = \xi_1, \end{cases} \quad (1)$$

where  $T > 0$ ,  $f : [a, T] \times \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}^m$  is a smooth function and  $\xi_0, \xi_1 \in \mathbb{R}^m$ . The existence and uniqueness of the solutions of (1) has been discussed in [13].

The mathematical modeling of physical phenomena in science and engineering often give rise to problems of this nature.

In order to avoid increasing the dimension of the problem, a situation associated with reducing (1) to a system of first order equations, (See [11]), we intend to solve the problem directly. This latter approach is preferred due to its advantages over the former, [4, 5].

In what follows, we shall develop an implicit single step method which is a hybrid of the continuous linear multistep method proposed in [4]. This we hope to achieve by the inclusion of offstep points. The motivation for the development of this method comes from the desire to combine the advantages in single step methods with the efficiency of multistep methods, (see [7, 8] and [12]).

The derivation and specification of the algorithm is presented in section two. In section three, we characterize the algorithm in terms of convergence and region of stability and then the implementation is discussed in section four. Lastly, numerical experiments are conducted to determine the efficiency of the algorithm

---

<sup>1</sup>Department of Mathematics, Covenant University, Canaanland, Ota, Ogun State, Nigeria,  
e-mail: timothy.anake@covenantuniversity.edu.ng, sheila.bishop@covenantuniversity.edu.ng,  
olasunmbo.agboola@covenantuniversity.edu.ng

*Manuscript received February 2015.*

## 2. DERIVATION OF THE METHOD

Let the polynomial of the form

$$y(x) = \sum_{i=0}^m a_i x^i \quad (2)$$

approximate the exact solution of (1) in the partition  $a = x_0 < x_1 < \dots < x_n < x_{n+1} < \dots < x_N = T$ , such that the step-size  $h$ , is defined as  $h = x_{n+1} - x_n$ ,  $n = 0, 1, \dots, N - 1$ .

Now, consider the one step design:  $x_n < x_{n+\nu_s} < x_{n+1}$ ,  $s = 1, \dots, t$ , where  $\nu_s$ ,  $\forall s$ , are offstep points and  $\nu \in \mathbb{Q}$ . Then, by interpolation of (2) at points  $x_{n+\nu_{t-1}}$ , and  $x_{n+\nu_t}$  and collocation of (1) at  $x_{n+r}$ ,  $r = 0, \nu_s, 1$ , we obtained the system

$$y_{n+\nu_s} = \sum_{i=0}^8 a_i x_{n+\nu_s}^i, \quad s = t - 1, t, \quad (3a)$$

$$f_{n+r} = \sum_{i=0}^8 i(i-1)a_i x_{n+r}^{i-2}, \quad r = 0, \nu_s, 1. \quad (3b)$$

Equations (3a) and (3b) are respectively interpolation and collocation equations. The system (3) is solved for the unknown coefficients,  $a_i$  which are substituted into (2). Using a scaling factor,  $z \in \mathbb{Z}$ , defined as  $z = \frac{x-x_{n+\nu_t}}{h}$ , such that  $z \leq 1$ , a continuous representation of our implicit hybrid algorithm and its first derivative is obtained as follows:

$$y_{n+r} = \Phi_{n+\nu}(z) + h^2 \left[ \sum_{i=0}^1 \beta_i(z) f_{n+i} + \sum_{s=1}^t \beta_{\nu_s}(z) f_{n+\nu_s} \right], \quad (4a)$$

$$\dot{y}_{n+r} = \dot{\Phi}_{n+\nu}(z) + h \left[ \sum_{i=0}^1 \dot{\beta}_i(z) f_{n+i} + \sum_{s=1}^t \dot{\beta}_{\nu_s}(z) f_{n+\nu_s} \right], \quad (4b)$$

where  $\Phi_{n+\nu}(z) = \alpha_{\nu_{(t-1)}}(z)x_{n+\nu_{(t-1)}} + \alpha_{\nu_{(t)}}(z)x_{n+\nu_{(t)}}$ . Obviously,  $\alpha_r(z)$  and  $\beta_r(z)$ , are continuous coefficients of  $z$ . By way of definition,  $y_{n+j} = y(x_{n+jh})$ , is the approximation of the exact solution at the point  $x_{n+j}$ ,  $f_{n+j} = f(x_{n+j}, y(x_{n+j}), \dot{y}(x_{n+j}))$ , for arbitrary  $j$ .

In particular, choosing  $\nu = \frac{s}{6}$ ,  $s = 1, \dots, 5$ , i.e  $t = 5$  the procedure described above yields the discrete schemes

$$y_{n+r} = \alpha_{\frac{2}{3}}(z)y_{n+\frac{2}{3}} + \alpha_{\frac{5}{6}}(z)y_{n+\frac{5}{6}} + h^2 \left[ \sum_{i=0}^1 \beta_i(z) f_{n+i} + \sum_{s=1}^5 \beta_{\nu_s}(z) f_{n+\nu_s} \right], \quad (5a)$$

$$\dot{y}_{n+r} = \dot{\alpha}_{\frac{2}{3}}(z)y_{n+\frac{2}{3}} + \dot{\alpha}_{\frac{5}{6}}(z)y_{n+\frac{5}{6}} + h \left[ \sum_{i=0}^1 \dot{\beta}_i(z) f_{n+i} + \sum_{s=1}^5 \dot{\beta}_{\nu_s}(z) f_{n+\nu_s} \right]. \quad (5b)$$

Evaluating (5a) for  $z = -\frac{5}{6}, -\frac{2}{3}, -\frac{1}{2}, -\frac{1}{3}$  and  $\frac{1}{6}$  yield discrete schemes whose coefficients are given in Table 1. Also, evaluating (5b) for  $z = -\frac{5}{6}, -\frac{2}{3}, -\frac{1}{2}, -\frac{1}{3}, -\frac{1}{6}, 0$  and  $\frac{1}{6}$  yield discrete derivative schemes described by their coefficients in Table 2.

Our main numerical algorithm and its first derivative are obtained as follows:

$$y_{n+1} = 2y_{n+\frac{5}{6}} - y_{n+\frac{2}{3}} + \frac{h^2}{2177280} \left[ 4315f_{n+1} + 53994f_{n+\frac{5}{6}} - 2307f_{n+\frac{2}{3}} \right. \\ \left. + 7948f_{n+\frac{1}{2}} - 4827f_{n+\frac{2}{3}} + 1578f_{n+\frac{1}{6}} - 221f_n \right], \quad (6)$$

$$y'_{n+1} = 6y_{n+\frac{5}{6}} - 6y_{n+\frac{2}{3}} + \frac{h}{241920} \left[ 12313f_{n+1} + 55246f_{n+\frac{5}{6}} - 18689f_{n+\frac{2}{3}} \right. \\ \left. + 19460f_{n+\frac{1}{2}} - 10921f_{n+\frac{2}{3}} + 3470f_{n+\frac{1}{6}} - 479f_n \right]. \quad (7)$$

TABLE 1. The coefficients for the method (5a)

r	$\alpha_{\frac{2}{3}}$	$\alpha_{\frac{5}{6}}$	$\beta_0$	$\beta_{\frac{1}{6}}$	$\beta_{\frac{1}{3}}$	$\beta_{\frac{1}{2}}$	$\beta_{\frac{2}{3}}$	$\beta_{\frac{5}{6}}$	$\beta_1$
0	5	-4	$\frac{409}{217728}$	$\frac{6366}{217728}$	$\frac{11679}{217728}$	$\frac{18532}{217728}$	$\frac{21255}{217728}$	$\frac{2334}{217728}$	$\frac{-95}{217728}$
$\frac{1}{6}$	4	-3	$\frac{-11}{362880}$	$\frac{758}{362880}$	$\frac{10503}{362880}$	$\frac{19708}{362880}$	$\frac{26883}{362880}$	$\frac{2754}{362880}$	$\frac{-95}{362880}$
$\frac{1}{3}$	3	-2	$\frac{31}{725760}$	$\frac{-207}{725760}$	$\frac{2313}{725760}$	$\frac{19708}{725760}$	$\frac{35073}{725760}$	$\frac{3762}{725760}$	$\frac{-137}{725760}$
$\frac{1}{2}$	2	-1	$\frac{31}{2177280}$	$\frac{-186}{2177280}$	$\frac{213}{2177280}$	$\frac{5426}{2177280}$	$\frac{49353}{2177280}$	$\frac{5862}{2177280}$	$\frac{-221}{2177280}$
1	-1	2	$\frac{-221}{2177280}$	$\frac{1578}{2177280}$	$\frac{-4827}{2177280}$	$\frac{7948}{2177280}$	$\frac{-2307}{2177280}$	$\frac{53994}{2177280}$	$\frac{4315}{2177280}$

TABLE 2. The coefficients for the method (5b)

r	$\alpha_{\frac{2}{3}}$	$\alpha_{\frac{5}{6}}$	$\beta_0$	$\beta_{\frac{1}{6}}$	$\beta_{\frac{1}{3}}$	$\beta_{\frac{1}{2}}$	$\beta_{\frac{2}{3}}$	$\beta_{\frac{5}{6}}$	$\beta_1$
0	-6	6	$\frac{-12287}{241920}$	$\frac{-58738}{241920}$	$\frac{-19697}{241920}$	$\frac{-58876}{241920}$	$\frac{-26465}{241920}$	$\frac{-6962}{241920}$	$\frac{585}{241920}$
$\frac{1}{6}$	-6	6	$\frac{1313}{725760}$	$\frac{-45990}{725760}$	$\frac{-149013}{725760}$	$\frac{-101620}{725760}$	$\frac{-119817}{725760}$	$\frac{-8262}{725760}$	$\frac{29}{725760}$
$\frac{1}{3}$	-6	6	$\frac{-413}{725760}$	$\frac{4266}{725760}$	$\frac{-55035}{725760}$	$\frac{-134132}{725760}$	$\frac{-105219}{725760}$	$\frac{-12438}{725760}$	$\frac{571}{725760}$
$\frac{1}{2}$	-6	6	$\frac{43}{241920}$	$\frac{-418}{241920}$	$\frac{2201}{241920}$	$\frac{-19708}{241920}$	$\frac{-39587}{241920}$	$\frac{-3074}{241920}$	$\frac{63}{241920}$
$\frac{2}{3}$	-6	6	$\frac{-253}{725760}$	$\frac{1962}{725760}$	$\frac{-6939}{725760}$	$\frac{15884}{725760}$	$\frac{-57123}{725760}$	$\frac{-14742}{725760}$	$\frac{731}{725760}$
$\frac{5}{6}$	-6	6	$\frac{289}{725760}$	$\frac{-2214}{725760}$	$\frac{7659}{725760}$	$\frac{-16628}{725760}$	$\frac{36855}{725760}$	$\frac{35514}{725760}$	$\frac{-995}{725760}$
1	-6	6	$\frac{-479}{241920}$	$\frac{3470}{241920}$	$\frac{-10921}{241920}$	$\frac{19460}{241920}$	$\frac{-18689}{241920}$	$\frac{55246}{241920}$	$\frac{12393}{241920}$

### 3. ANALYSIS OF THE METHODS

In this section, we set out to test the convergence and determine the region of stability of the new algorithm, (6) .

3.1. **Order and error constant.** We begin by rewriting (6) as a linear difference operator of the form:

$$\mathcal{L}[y(x); h] = y(x) - \alpha_{\nu_{t-1}}y(x_n + \nu_{(t-1)}h) - \alpha_{\nu_n}y(x_n + \nu_t h) - h^2 \left[ \sum_{i=0}^1 \beta_i \ddot{y}(x_n + ih) + \sum_{s=1}^t \beta_{\nu_s} \ddot{y}(x_n + \nu_s h) \right], \tag{8}$$

where  $y(x) \in C^d[a, T]$ ,  $d \in \mathbb{R}$ , is an arbitrary test function. Then expand  $y(x_n + rh)$  and  $\ddot{y}(x_n + rh)$ ,  $r = 0, \nu_u, 1$  for all  $r$  respectively in Taylor series about  $x_n$  and collecting terms in powers of  $y$  gives

$$\mathcal{L}[y(x); h] = C_0y(x) + C_1hy^{(1)}(x) + C_2h^2y^{(2)}(x) + \dots + C_ph^py^{(p)}(x) + C_{p+1}h^{p+1}y^{(p+1)}(x) + \dots, \tag{9}$$

where the constant coefficients  $C_q, q = 0, 1, 2, \dots$  are defined as follows:

$$\begin{aligned}
 C_0 &= \sum_{i=0}^k \alpha_i \\
 C_1 &= \sum_{i=0}^k i\alpha_i \\
 &\vdots \\
 C_q &= \frac{1}{q!} \left[ \sum_{i=1}^k i^q \alpha_i - q(q-1) \left( \sum_{i=1}^k i^{q-2} \beta_i + \sum_{i=1}^k \mu_i^{(q-2)} \beta_{\mu_i} \right) \right].
 \end{aligned}
 \tag{10}$$

**Definition 3.1.** (1) The difference operator  $\mathcal{L}$  and the associated method is said to be of order  $p$  if  $C_0 = C_1 = \dots = C_p = 0$  and  $C_{p+2} \neq 0$   
 (2) The term  $C_{p+2}$  is called the error constant and it implies that the local truncation error (l.t.e) is defined by:

$$l.t.e = C_{p+2} h^{p+2} y^{p+2} + O(h^{p+3}). \tag{11}$$

We have established from our computation that methods (6) has order  $p = 7$  and error constants  $C_{p+2} = -3.1173 \times 10^{-10}$ .

**3.2. Zero stability and consistency.**

**Definition 3.2.** The method (6) is said to be zero stable as  $h \rightarrow 0$  if its first characteristic polynomial  $\rho(z)$ , satisfies the condition  $|z_\phi| \leq 1$ . Furthermore, those roots for which  $|z_\phi| = 1$  have multiplicity not exceeding 2.

The first characteristic polynomial of the algorithm is obtained as follows;

$$\rho(z) = z - 2z^{5/6} + z^{2/3} = 0. \tag{12}$$

It can be seen that the conditions in Definition (3.2) are satisfied by (12). Therefore, we conclude that the numerical algorithm (6) is zero stable. The consistency of the method is established also, by the fact that the order  $p = 7$  of the algorithm is greater than 1, that is,  $\check{\rho}(1) = 2!\sigma(1)$  (see [10]).

**3.3. Convergence.** Since we have established the consistency and zero stability of the algorithm, we can conclude that the numerical algorithm (6) is convergent (see [9]).

**3.4. Region of absolute stability.** Absolute stability for the algorithms is determined by means of the boundary locus method. Consider the stability polynomial

$$\Pi(z, \bar{h}) = \rho(z) - \bar{h}\sigma(z) = 0, \tag{13}$$

where  $\bar{h} = h^2 \lambda^2$  and  $\lambda = \frac{df}{dy}$  are assumed constant.

The polynomial (13) is obtained by applying the continuous implicit one step hybrid algorithm (6) to the scalar test problem;

$$y'' = -\lambda^2 y. \tag{14}$$

**Definition 3.3.** The algorithm, (6) is said to be absolutely stable if for a given  $\bar{h}$  all the roots  $z_\phi$  of (13) satisfy  $|z_\phi| < 1, \phi = 1, 2, \dots, (r - 1)$ .

**Definition 3.4.** The region  $\mathcal{R}$  of the complex  $\bar{h}$ -plane such that the roots of the polynomial  $\Pi(z, \bar{h})$  lie within the unit circle whenever  $\bar{h}$  lies in the interior of the region is called the region of absolute stability.

We established from our computation using the boundary locus method, that our algorithm, (6) has interval of absolute stability to be  $(-9.87, 0)$ .

**3.5. Implementation of the method.** The method (6), we use simultaneous solutions of a zero stable block formula of the form (the zero stability of the block formula has been established in [3]). :

$$h^\gamma \mathbf{A}Y_m^{(n)} = h^\gamma \mathbf{B}y_m^{(n)} + h^{\mu-\gamma} [\mathbf{C}\mathbf{F}(Y_m)], \tag{15}$$

where  $Y_m^{(n)} = (y_{n+\nu_1}, \dots, y_{n+\nu_{(t-1)}}, y_{n+\nu_{(t)}}, y_{n+1}, \dot{y}_{n+\nu_1}, \dots, \dot{y}_{n+\nu_{(t-1)}}, \dot{y}_{n+\nu_{(t)}}, \dot{y}_{n+1})^T$ ;  
 $y_m^{(n)} = (y_{n-\nu_1}, \dots, y_{n-\nu_{(t-1)}}, y_{n-\nu_{(t)}}, y_n, \dot{y}_{n-\nu_1}, \dots, \dot{y}_{n-\nu_{(t-1)}}, \dot{y}_{n-\nu_{(t)}}, \dot{y}_n)^T$ ;  
 $\mathbf{F}(Y_m) = (f_n, f_{n+\nu_1}, \dots, f_{n+\nu_{(t-1)}}, f_{n+\nu_{(t)}}, f_{n+1})^T$ ;  $n$  represents the order of the derivative of (6);  $\gamma = n$  is the power of  $h$ ;  $\mu$  is the order of (1);  $\mathbf{A}$ ,  $\mathbf{B}$  and  $\mathbf{C}$  are constant coefficient matrices obtained as follows:  $\mathbf{A}$  is  $12 \times 12$  identity matrix,

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ \frac{1}{6} & \frac{1}{3} & \frac{1}{2} & \frac{2}{3} & \frac{5}{6} & 1 & 1 & 1 & 1 & 1 \end{bmatrix}^T$$

and

$$C = \begin{bmatrix} \frac{28549}{4354560} & \frac{275}{20736} & -\frac{5717}{483840} & \frac{10621}{1088640} & -\frac{7703}{1451520} & \frac{403}{241920} & -\frac{199}{870912} \\ \frac{1027}{68040} & \frac{97}{1890} & -\frac{2}{81} & \frac{197}{8505} & \frac{97}{7560} & \frac{23}{5670} & -\frac{19}{34120} \\ \frac{253}{10752} & \frac{165}{1792} & -\frac{267}{17920} & \frac{5}{128} & -\frac{363}{17920} & \frac{57}{8960} & -\frac{47}{53760} \\ \frac{272}{8505} & \frac{376}{2835} & -\frac{2}{945} & \frac{656}{8505} & -\frac{2}{81} & \frac{8}{945} & -\frac{2}{1701} \\ \frac{35225}{870912} & \frac{8375}{48384} & \frac{3125}{290304} & \frac{25625}{217728} & -\frac{625}{96768} & \frac{275}{20736} & -\frac{1375}{870912} \\ \frac{41}{840} & \frac{3}{14} & \frac{3}{140} & \frac{17}{105} & \frac{3}{280} & \frac{3}{70} & 0 \\ \frac{19087}{362880} & \frac{2703}{15120} & -\frac{15487}{120960} & \frac{293}{2835} & -\frac{6737}{120960} & \frac{263}{15120} & -\frac{863}{362880} \\ \frac{1139}{22680} & \frac{47}{189} & \frac{11}{7560} & \frac{166}{2835} & -\frac{269}{7560} & \frac{11}{945} & -\frac{37}{22680} \\ \frac{137}{2688} & \frac{27}{112} & \frac{387}{4480} & \frac{17}{105} & -\frac{243}{4480} & \frac{9}{560} & -\frac{29}{13440} \\ \frac{143}{2835} & \frac{232}{945} & \frac{64}{945} & \frac{752}{2835} & \frac{29}{945} & \frac{8}{945} & -\frac{4}{2835} \\ \frac{3715}{7257} & \frac{725}{3024} & \frac{2125}{24192} & \frac{125}{567} & \frac{3875}{24192} & \frac{235}{3024} & -\frac{275}{72576} \\ \frac{41}{840} & \frac{9}{25} & \frac{9}{280} & \frac{34}{105} & \frac{9}{280} & \frac{9}{35} & \frac{41}{840} \end{bmatrix}$$

This is a block by block procedure, where starting values for subsequent blocks are obtained from the previous block for the implementation of the method over the subintervals:  $[x_0, x_1], [x_1, x_2], \dots, [x_{N-1}, x_N]$ .

#### 4. NUMERICAL EXPERIMENT

The derived method is experimented on some sample problem to illustrate its performance. Our results are compared with previous results obtained from other authors.

**Problem 4.1.**  $y'' - x(y')^2 = 0, y(0) = 1, y'(0) = \frac{1}{2}; h = 0.01.$

Theoretical solution:

$$y = 1 + \frac{1}{2} \ln \left( \frac{2+x}{2-x} \right).$$

**Problem 4.2.**  $y'' = \frac{(y')}{2y} - 2y, y \left( \frac{\pi}{6} \right) = \frac{1}{4}, y' \left( \frac{\pi}{6} \right) = \frac{\sqrt{3}}{2}; h = 0.01.$

Theoretical Solution:

$$y(x) = (\sin x)^2.$$

TABLE 3. Absolute errors in the new method to errors in [1, 2] for Problem 4.1

X	Exact Result	Computed Result	Error in method (6)	Error in [1]	Error in [2]
0.1	1.050041728	1.050041728	6.2172E-15	7.5028(-13)	4.8627(-14)
0.2	1.100335345	1.100335345	2.4425E-14	9.7410(-12)	2.1604(-13)
0.3	1.151140433	1.151140433	5.6843E-14	3.7638(-11)	5.2557(-13)
0.4	1.202732549	1.202732549	1.0347E-13	9.7765(-11)	1.0254(-12)
0.5	1.255412806	1.255412806	1.6742E-13	2.0825(-10)	1.8032(-12)
0.6	1.309519597	1.309519597	2.5091E-13	3.9604(-10)	3.0078(-12)
0.7	1.365443745	1.365443745	3.6016E-13	7.0460(-10)	4.8991(-12)
0.8	1.423648920	1.423648920	5.0493E-13	1.2095(-09)	7.9460(-12)
0.9	1.484700266	1.484700266	6.9522E-13	2.0511(-09)	1.3702(-11)
1.0	1.549306129	1.549306129	9.4836E-13	3.5066(-09)	2.1885(-11)

TABLE 4. Absolute errors in method (6) to errors in [1, 6] for Problem 4.2

X	Exact Result	Computed Result	Error in Method (6)	Error in [1]	Error in [6]
1.1035988	0.7971525560	0.7971525560	3.9946E-11	1.8811(-10)	2.8047(-10)
1.2035988	0.8711181669	0.8711181669	3.3286E-11	2.4539(-10)	2.7950(-10)
1.3035988	0.9302884674	0.9302884673	2.5298E-11	3.0306(-10)	2.1490(-10)
1.4035988	0.9723045242	0.9723045242	1.6302E-11	3.5819(-10)	5.4975(-11)
1.5035988	0.9954912899	0.9954912899	6.6553E-12	4.0838(-10)	1.1545(-10)
1.6035988	0.9989243811	0.9989243811	3.2564E-12	4.5128(-10)	4.4825(-10)
1.7035988	0.9824669315	0.9824669315	1.3038E-11	5.0696(-10)	1.1840(-09)
1.8035988	0.9467750474	0.9467750475	2.2300E-11	5.0696(-10)	1.1840(-09)
1.9035988	0.8932716519	0.8932716519	3.0673E-11	5.1697(-10)	1.6318(-09)
2.0035988	0.8240897564	0.8240897565	3.7823E-11	5.1381(-10)	2.0567(-09)

## 5. CONCLUSION

Hybrid numerical algorithm of a one step method of order  $p = 7$  has been developed in this paper. The algorithm gives convergent solutions with very low errors and is absolutely stable. Implementation of the algorithm is by block method which apart from generating simultaneous solutions also gives derivative solutions. The experiment performed with the numerical algorithm proposed in this paper shows that it performs better than the methods proposed in earlier [1, 2] and [6].

## REFERENCES

- [1] Adesanya, A.O., Odekunle, M.R., Adeyeye, A.O., (2012), Continuous block hybrid-predictor-corrector method for the solution of  $y'' = f(x, y, y')$ , Intern. J. of Math. and Soft Computing, 2(2), pp.35-42.
- [2] Adesanya, A.O., Odekunle, M.R., Alkali, A.A., (2012), Order six block predictor-corrector method for the solution of  $y'' = f(x, y, y')$ , Canadian J. of Sci. and Engr., 3(4), pp.180-185.
- [3] Anake, T.A., (2011), Continuous Implicit Hybrid One Step Methods for the Solutions of Initial Value Problems of General Second Order Ordinary Differential Equations, Ph.D. Thesis, Covenant University, Canaanland, Ota, Ogun State, Nigeria.
- [4] Awoyemi, D.O., (1992), On some Continuous Linear Multistep Methods for Initial Value Problems, Ph.D. Thesis, University of Ilorin, Ilorin, Nigeria.
- [5] Awoyemi, D.O., (2001), A new Sixth-order algorithm for general secondary order ordinary differential equation, Intern. J. Comput. Math, 77, pp.117-124.

- [6] Awoyemi, D.O., Adebile, E.A., Adesanya, A.O., Anake, T.A., (2011), Modified block method for the direct solution of second order ordinary differential equations, *Intern. J. of Appl. Math. and Comput.*, 3(3), pp.181-188.
  - [7] Brush, D.G., Kohfield, J.J., Thompson, G.T., (1967), Solution of ordinary differential equations using two “off-step” points, *Journal of the ACM*, 14(4), pp.769-784.
  - [8] Gear, C.W., (1964), Hybrid methods for initial value problems in ordinary differential equations, *SIAM Journal of Numerical Analysis*, 2, pp.69-86.
  - [9] Henrici, P., (1962), *Discrete Variable Methods in ODE*, John Wiley and Sons, New York.
  - [10] Lambert, J.D., (1973), *Computational Methods in Ordinary Differential Equations*, John Wiley, New York, p.278.
  - [11] Shampine, L.F., (1994), *Numerical Solution of Ordinary Differential Equations*, Chapman and Hall, New York, p.484.
  - [12] Stanimirovic, Z., Maric, M., Radijicic N., Bozovic S., (2014), Two efficient hybrid metaheuristic methods for solving the load balance problem, *Appl. Comput. Math.*, 13(3), pp.332-349.
  - [13] Wend, D.V., (1969), Existence and uniqueness of solutions of ordinary differential equations, *Proceedings of the American Mathematical Society*, 23(1), pp.27-33.
- 
- 



**Timothy A. Anake** is currently a Lecturer and Postgraduate Studies Coordinator of the Department of Mathematics, School of Natural and Applied Sciences, College of Science and Technology, Covenant University, Canaanland. He got B.Sc. degree in Mathematics from the University of Uyo and M.Sc. degree in Mathematics from the University of Ibadan. In 2011 he got Ph.D. degree in Mathematics from Covenant University, Ota. His current research interests include numerical solutions of differential equations, moving boundary problems using mesh free methods and solutions of Schrodinger equations.



**Sheila A. Bishop** graduated from the University of Benin in 1998 with a Bachelor of Science Education degree in Mathematics and obtained a Master degree in Mathematics from the University of Ibadan in 2005. She joined Covenant University in 2007 as an assistant Lecturer in the Department of Mathematics. She obtained her Ph.D. degree in Mathematics in 2012. She has a teaching experience of over a decade in both public and private schools.



**Olasunmbo O. Agboola** holds Bachelor of Science Education (First Class Hons.) and Master of Science degrees from the University of Ilorin, Ilorin, Nigeria. He is an Assistant Lecturer in the Department of Mathematics in the College of Science and Technology. His research interests are in analytical dynamics and numerical analysis. Presently, he is undertaking his Doctor of Philosophy (Ph.D) degree programme at Covenant University where he equally teaches undergraduate courses in Mathematics.